Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 1 February 2008 (MN LATEX style file v1.4)

The infrared luminosity of the torus and the visibility of scattered broad line emission in Seyfert 2 galaxies

S.L. Lumsden[1] and D.M. Alexander[2]

[1] Department of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, UK
Email – sll@ast.leeds.ac.uk
[2] Department of Astronomy & Astrophysics, 525 Davey Laboratory, Pennsylvania State University, University Park, PA 16802, USA
Email – davo@astro.psu.edu

arXiv:astro-ph/0110174v1 8 Oct 2001

1 February 2008

ABSTRACT

A number of studies have shown that the visibility of scattered broad emission lines in Seyfert 2 galaxies is strongly dependent on the IRAS f_{60}/f_{25} flux ratio, where those Seyfert 2s with "warm" IRAS colours show polarised broad line emission. It is now clear that this effect is due to the increasing dominance of the galactic rather than the AGN emission at 60 μm in less luminous "cool" Seyfert 2s. However, we present evidence that the 25 μm emission is a good measure of the AGN luminosity for most Seyfert 2s. Using this result, we show that the visibility of scattered broad line emission has a dependence on the AGN luminosity. The observations can be interpreted self-consistently if the scale height of the scattering zone varies with central source luminosity whilst the scale height of the obscuring torus is approximately constant.

Key words: galaxies: Seyfert - galaxies: active - polarization - scattering - infrared: galaxies - X-rays: galaxies

1 INTRODUCTION

The unified model for Seyfert galaxies provides a good qualitative explanation of the apparently diverse properties of Seyfert 1 and Seyfert 2 galaxies (e.g., Antonucci 1993). In this model, both types of Seyfert have the same nuclear components, namely a broad line region and accretion disc around a central supermassive black hole, but in Seyfert 2 galaxies the nucleus is obscured by an optically and geometrically thick dusty torus. Strong support for the dusty torus model has come from both theoretical models (e.g. Pier & Krolik 1993; Granato & Danese 1994; Efstathiou & Rowan-Robinson 1995) and X-ray and near-IR observations (e.g., Alonso-Herrero, Ward, & Kotilainen 1997; Turner et al. 1997; Risaliti, Maiolino, & Salvati 1999). Direct evidence for hidden Seyfert 1 activity in Seyfert 2 galaxies has come from optical spectropolarimetric observations which have shown that broad line emission is observed in scattered flux in some Seyfert 2 galaxies (e.g., Antonucci & Miller 1985; Young et al. 1996; Heisler, Lumsden, & Bailey 1997 hereafter HLB; Moran et al. 2000; Lumsden et al. 2001 hereafter L01; Tran 2001).

HLB carried out the first statistical study of the frequency of polarised broad line emission in Seyfert 2 galaxies and found a correlation between the occurrence of polarised broad line emission and the IRAS f_{60}/f_{25} flux ratio where only those Seyfert 2s with "warm" IRAS colours (f_{60}/f_{25} <4.0) showed polarised broad line emission. It was suggested in that paper that the IRAS colours reflected the the inclination of the torus, so that "cool" sources had more obscured nuclei. However, this explanation was later shown to be inconsistent with the evidence provided from X-ray observations (Alexander 2001; hereafter A01). Instead, A01 proposed an alternative picture where the IRAS f_{60}/f_{25} flux ratio was related to the relative luminosities of galactic and AGN activity. In this picture the lack of polarised broad line emission in "cool" Seyfert 2 galaxies is due to either an intrinsically weak AGN component or a dominant galactic component. Further analysis by Gu et al. (2001) and L01 supported this picture and it is now clear that the f_{60}/f_{25} flux ratio is not directly related to the inclination of the torus in the large IRAS apertures.

However, the previous data suggested that the 25μm emission was related to the AGN, and the primary aim of this paper is to test that result by examining the correlation between the 25 μm luminosity (hereafter L_{25}) and extinction corrected hard X-ray (HX, 2–10keV) and [OIII] 5007Å luminosities (hereafter L_{HX} and L_{5007}). We use these results to test how the detectability of polarised broad lines in Seyfert 2s depends on AGN luminosity, since this has never been properly examined in the past, and suggest a scheme that explains our results.

© 0000 RAS

2 MID INFRARED EMISSION IN SEYFERT GALAXIES

The most commonly used indicators of the AGN luminosity are L_{HX} and L_{5007} (eg. Mulchaey et al. 1994, Alonso-Herrero et al. 1997). Unfortunately there is no single homogeneous sample of Seyfert galaxies observed in either band. At the time of writing the most complete compilation of Seyfert 2 HX properties is presented by Bassani et al. (1999) and we have used this sample for our HX analysis. To provide a comparison, we have included the Seyfert 1 galaxies from Reynolds (1997). The heterogeneous nature of these X-ray samples is not in itself a problem, since the observed galaxies span a wide range of obscuration and luminosity, which are the primary parameters that we might expect to introduce a bias in any correlations we find. However a potential bias could be introduced by the nature of the observations. Unlike the infrared or optical data, the X-ray catalogue is essentially flux limited (where the limit is set by the available instrument/mission sensitivity). This could translate into a false luminosity correlation if there are a large number of galaxies at the flux limit, but with widely varying redshifts. We have attempted to correct for this by only using the galaxies with $cz < 9000\mathrm{kms}^{-1}$. We have also excluded all sources from both samples which are radio-loud or have optical characteristics more typical of LINERs or quasars so that the sample does at least have a homogeneous optical classification. The X-ray emission was corrected for absorption in the case of Compton thin sources (those with $N_H < 1.5 \times 10^{24}\mathrm{cm}^{-2}$). No correction was made for the Compton thick sources since the direct X-ray emission is blocked in this case and the detected emission is predominantly scattered light. Estimating L_{HX} in these sources is dependent on the scattering model adopted, and hence prone to error; we have not included these galaxies in the correlation tests involving L_{HX}.

The [OIII] 5007Å data was taken from Bassani et al. for the Seyfert 2s, and from Mulchaey et al. (1994) for the Seyfert 1s. The [OIII] fluxes given by Bassani et al. are already extinction corrected: the data for the Seyfert 1s was corrected using, in order of preference, the Hα/Hβ ratios given in Mulchaey et al. (1994) and the values of A_V listed in Winkler et al. (1992). We discarded those galaxies without reported [OIII] fluxes. IRAS 25μm fluxes for all these galaxies were taken from, in order of preference, the IRAS Bright Galaxy Survey (Soifer et al. 1989, Sanders et al. 1995), the 1.2 Jy catalog (Strauss et al. 1990) and the IRAS Faint Source Catalogue (Moshir et al. 1991). We define $L_{25} = \nu P_\nu(25\mu\mathrm{m})$, where P_ν is the radiant power. The IRAS data was not corrected for any residual opacity at this wavelength. We discarded three galaxies which either lie in gaps in the IRAS sky coverage (MCG-5-23-16 and NGC3081) or are blended with other nearby bright sources (NGC2992 where the IRAS emission is a blend of the emission from NGC2992 and NGC2993). We used the survival analysis techniques in the software package ASURV (version 1.1: La Valley, Isobe & Feigelson 1992) to deal with the upper limits present in the IRAS data. This implements the methods for correlation analysis outlined in Isobe, Feigelson & Nelson (1986).

Figure 1 shows the correlation between L_{25} and (a) L_{HX} and (b) L_{5007}. Correlations between the three are found at $>99\%$ confidence. We checked for any bias introduced by the essentially flux limited X-ray samples by also examining the correlations between the fluxes. Correlations of the [OIII] 5007Å flux with both the 25μm and HX fluxes are also found with $>99\%$ confidence. The correlation of the 25μm and HX fluxes is weaker, with only 93% confidence. There is no evidence from this however for any bias being introduced into the luminosity correlations we have found.

There are trends in the luminosity correlations that the raw probabilities do not reflect. The galaxies with lower L_{HX} or L_{5007} tend to have excess 25μm emission. It is likely that there is a significant contribution from the host galaxy emission (presumably star formation) in these galaxies. We tested for the presence of an additional host galaxy component in two ways. First, we examined the IRAS colours of the galaxies in the fashion of Dopita et al. (1998). Figure 2(a) shows that objects with low L_{5007} clearly have IRAS colours typical of starbursts or LINERs (see also Figure 1 in L01). Secondly, we examined the compactness of the 10μm emission as a function of L_{5007}, using the data from Maiolino et al. (1995) and Giuricin, Mardirossian and Mezzetti (1995). The compactness is defined in the standard sense as the colour-corrected ratio of the small aperture (3–10$''$) ground based 10μm flux and the IRAS 12 μm flux. We assume that the 10 and 25μm emission trace the same dust (it would be better to measure the compactness at 20μm but there are very few small aperture observations available). Figure 2(b) shows that there is a strong correlation between the compactness and the AGN luminosity at >99.9% confidence for galaxies with $cz < 5000\mathrm{kms}^{-1}$. We truncated the redshift range in order to avoid false correlations caused by the fact that the more luminous systems also tend to be more distant. The reality of this correlation is demonstrated by the fact that the actual observed confidence level does not change if we truncate the sample anywhere in the range $2500 < cz < 25000\mathrm{kms}^{-1}$. There is clear evidence from both these indicators that the host galaxy can be a significant contributor at these wavelengths in less luminous Seyferts.

We estimated the luminosity at which galactic 25μm emission has a significant effect on the overall spectral energy distribution by inspection of Figure 2(a). We found that galaxies with $L_{HX} \lesssim 10^{7.2}\,L_\odot$ and $L_{5007} \lesssim 10^{7.3}\,L_\odot$ were likely to have IRAS colours more typical of the host galaxy. We therefore excluded these systems when deriving the fits to the data shown in Figure 1. Note however that the actual significance of the correlations seen in Figure 1 hardly changes if we exclude these lower luminosity AGN. We fitted the data with the bisector of the linear regression of y on x and x on y, which is the best estimate of the underlying trend in this case (see Isobe et al. 1990) The slopes found are all close to unity, within the errors, as expected if all three observational parameters reflect the underlying AGN luminosity. The actual values are 0.82±0.12 and 1.05±0.12 for the data shown in Figure 1(a) and (b) respectively, and 0.98±0.08 for the correlation between L_{HX} and L_{5007} (not shown in Figure 1, since it essentially repeats the previous work of Mulchaey et al. 1994, and Alonso-Herrero et al. 1997). If we include the galaxies with excess emission at 25μm the slope decreases to 0.72±0.06 in both Figure 1(a) and (b). This is in the sense expected if the lower luminosity galaxies do have excess 25μm emission. It is clear from these results that the L_{25} is as good an indicator of the underlying AGN luminosity as the extinction corrected X-ray or [OIII] luminosities, at least for those galaxies with $L_{5007} > 10^{7.3}\,\mathrm{L}_{\odot}$.

3 DETECTABILITY OF SCATTERED BROAD LINE EMISSION IN SEYFERT 2 GALAXIES

Scattered broad Hα from the broad line region has been detected in > 30 Seyfert 2s using optical spectropolarimetry since the initial observations of NGC1068 by Antonucci & Miller (1985). Until the work of HLB there were no statistically significant samples to test the mechanisms that effect the visibility of scattered broad line emission in Seyfert 2 galaxies. Now with the addition of the larger distance limited sample of Moran et al. (2000), the far infrared flux and luminosity limited sample of L01 and the heterogeneous optical and mid infrared selected sample of Tran (2001) we have a much firmer grasp of how often we see such *hidden* broad line regions (HBLRs). However, there have been no direct attempts to relate their detectability directly to the AGN luminosity, partly because suitable measures of that luminosity were not always available. Therefore, we have considered this issue using L_{25} as an indicator of the AGN luminosity using the above samples; we have implicitly assumed that all of the Seyfert 2s in the Moran et al. (2000) sample are not HBLRs unless they are included in their reported detections or were previously known to contain HBLRs (though we have included the two extra detections given in Moran et al. 2001). We have used the IRAS fluxes from the same sources as in Section 2. Some of the Moran et al. sample lie within gaps in the IRAS sky coverage and were excluded (MCG-5-23-16, NGC3081, NGC4117 and NGC7450), as was NGC2992 for the reason outlined in Section 2. Finally, we note that some of the galaxies in the Moran et al. sample only have upper limits at 25μm, necessitating the use of the ASURV package when comparing samples on the basis of luminosity.

We tested whether the redshift distribution of the HBLRs and non-HBLRs are the same for all three samples before comparing the luminosities to avoid introducing a Malmquist type bias into our analysis. A Mann-Whitney U test shows there is no evidence for a difference in the redshift distributions of HBLRs and non-HBLRs in the Moran et al. (2000) and L01 samples at the 5% significance level. The Tran (2001) sample does however show a difference at the 1% significance level. This is largely due to the presence of many low luminosity nearby AGN in the 12μm galaxy sample of Rush et al. (1993) which makes up well over 50% of the Tran sample. If we impose a lower redshift limit on the Tran sample of 2500kms^{-1}, then the difference in the redshift distribution is removed. Therefore in what follows we have used this truncated version of the Tran sample which has 2 HBLRs and 8 non-HBLRs removed. In practice, this has little effect on the actual significance of the difference in luminosities between the subsets found below, but is the simplest way to correct for this bias in the Tran sample. We also note that we have excluded the 2 galaxies from the L01 sample which had insufficient signal-to-noise to adequately determine the absence of an HBLR (see Section 3.1 of L01 for further details).

Figure 3 (a-c) shows the L_{25} distributions for the samples. Statistically the distributions for the HBLR samples differ from the non-HBLR samples at the 99level for both the Moran et al. and Tran samples (Figure 3a and 3b), with the galaxies containing HBLRs being more luminous. We used the four available two sample tests in ASURV to test whether the upper limits present affected the result: the statistical significance was $> 99\%$ for all four. These tests are outlined in Feigelson & Nelson (1985). There are fewer low luminosity Seyferts in the L01 sample, so the clear distinction present in the other samples is less obvious in Figure 3(c). Statistically the galaxies with and without HBLRs do have different L_{25} at the 88% confidence level, but this cannot be described as conclusive. We therefore also considered L_{5007}, which was available for this sample. The two samples differ here at the 92% confidence level in good agreement with the results of the 25μm data. Again this is not conclusive, but the similarity of the two indicators gives added confidence to our results for the Moran et al. and Tran samples, and the actual result agrees with the trends seen in those samples.

Finally we considered how our results are affected by our caveat that L_{25} may not truly reflect the AGN luminosity in the lowest luminosity galaxies. Clearly, all this can do is move some of the low luminosity galaxies (which uniformly show no evidence of an HBLR) to higher luminosities. Therefore the actual distinction between galaxies containing an HBLR and those without may be greater than we have estimated.

4 DISCUSSION AND CONCLUSIONS

We have shown that the large beam 25μm emission measured by IRAS is a good indicator of the underlying AGN luminosity in Seyfert galaxies for even the most obscured systems. Within the Unified model of Seyfert galaxies, any AGN emission at 25μm is produced by dust emission within the torus. Our results therefore suggest a direct relationship between the luminosity of the central source and the luminosity of the dusty torus, as was previously found for shorter wavelength infrared data by Alonso-Herrero et al. (1997). Furthermore, those sources with powerful AGN have warm IRAS f_{60}/f_{25} colours. A corollary of this and previous results of HLB is that those sources with luminous 25μm emission should therefore be more likely to have detectable HBLRs; we have directly shown this is the case for 3 independent optical spectropolarimetric samples.

An obvious conclusion for the lack of a HBLR could be that they do not exist in the lower luminosity Seyfert 2 galaxies (ie these are “true” Seyfert 2s). L01 discussed this issue, where it was shown that there is no significant difference in the detection rates of obscured HX emission or compact cm radio emission in galaxies with and without HBLRs. A similar result is shown directly by the HX study of Moran et al. (2001) and can also be inferred for the Moran et al. (2000) sample from the analysis of Gu et al. (2001). Clearly both galaxies with and without HBLRs show evidence for obscured or scattered Seyfert 1 cores in HX spectra.

How can the luminosity of the AGN affect the “visibility” of the HBLR? One possibility, addressed in detail in A01 and L01, is that the relatively larger contribution from the host galaxy in the less luminous AGN leads to increasing dilution of the polarisation signal, making it harder to detect any HBLR. Alternatively, we might be seeing a more direct physical connection between the AGN luminosity and the visibility of the HBLR. The scale height of the scattering particles will be larger in the more luminous sources for any reasonable geometry and electron density distribution. This creates two biases against the detection of HBLRs in less luminous sources. First, any HBLR emission will be intrinsically weaker (e.g., see A01) and secondly, the angular size of the scattering screens will be smaller. A smaller scattering screen has a larger probability of being obscured from view by either dust in the host galaxy or dust within the torus. The importance of the latter effect will depend upon the scale height of the torus and the relationship this has with the luminosity of the central source. There are several processes that could effect the scale height of the torus (e.g., radiation pressure will inflate the torus, gravitational and dynamical effects will tend to compress it). However, there are no forces that will tend to compress the size of the scattering volume. Therefore, the dependence on luminosity of the scale height of the scattering particles should be greater than that of the torus. The fraction of scattering that is obscured by the torus will therefore decrease as the AGN luminosity increases, naturally giving rise to the observed behaviour found in Section 3.

ACKNOWLEDGEMENTS

SLL acknowledges support from PPARC through the award of an Advanced Research Fellowship. DMA acknowledges financial support from the NSF CAREER award AST-9983783 (P.I. W.N.Brandt). We thank the referee Ed Moran for his very helpful comments.

REFERENCES

Alexander, D.M., 2001, MNRAS, 320, L15 (A01)
Alonso-Herrero, A., Ward, M.J., Kotilainen, J.K., 1997, MNRAS, 288, 977
Antonucci, R., Miller, J.S. 1985, ApJ, 297, 621
Antonucci, R. 1993, ARA&A, 31, 473
Bassani, L., Dadina, M., Maiolino, R., Salvati, M., Risaliti, G., della Ceca, R., Matt, G., Zamorani, G., 1999, ApJS, 121, 473
Dopita, M.A., Heisler, C.A., Lumsden, S.L., Bailey, J.A., 1998, ApJ, 498, 570
Efstathiou, A., Rowan-Robinson, M. 1995, MNRAS, 273, 649
Feigelson, E.D., Nelson, P.I., 1985, ApJ, 293, 192
Giuricin, G., Mardirossian, F., Mezzetti, M. 1995, ApJ, 438, 527
Granato, G.L., Danese, L. 1994, MNRAS, 268, 235
Gu, Q., Maiolino, R., Dultzin-Hacyan, D., 2001, A&A, 366, 765
Heisler, C.A., Lumsden, S.L., Bailey, J.A., 1997, Nature, 385, 700 (HLB)
Isobe, T., Feigelson, E.D., Nelson, P.I., 1986, ApJ, 306, 490
Isobe, T., Feigelson, E.D., Akritas, M.G., Babu, G.J., 1990, ApJ, 364, 104
LaValley, M., Isobe, T., Feigelson, E.D., 1992, BAAS, 24, 839
Lumsden, S.L., Heisler, C.A., Bailey, J.A., Hough, J.H., Young, S., 2001, MNRAS, in press (L01)
Maiolino, R., Ruiz, M., Rieke, G.H., Keller, L.D., 1995, ApJ, 446, 561
Moran, E.C., Barth, A.J., Kay, L.E., Filippenko, A.V., 2000, ApJ, 540, L73
Moran, E.C., Kay, L.E., Davis, M., Filippenko, A.V., Barth, A.J., 2001, ApJ, 556, L75
Moshir, M., et al., 1991, Explanatory Supplement to the IRAS Faint Source Survey, Version 2. JPL, Pasadena
Mulchaey, J.S., Koratkar, A., Ward, M.J., Wilson, A.S., Whittle, M., Antonucci, R.R.J., Kinney, A.L., Hurt, T., 1994, ApJ, 436, 586
Pier, E., Krolik, J. 1993, ApJ, 418, 673
Reynolds, C.S., 1997, MNRAS286, 513
Risaliti, G., Maiolino, R., Salvati, M. 1999, ApJ, 522, 157
Rush, B., Malkan, M.A., Spinoglio, L. 1993, ApJS, 89, 1
Sanders, D.B., Egami, E., Lipari, S., Mirabel, I.F., Soifer, B.T., 1995, AJ, 110, 1993
Soifer, B.T., Boehmer, L., Neugebauer, G., Sanders, D.B., 1989, AJ, 98, 766
Strauss, M.A., Davis, M., Yahil, A., Huchra, J.P., 1990, ApJ, 361, 49
Tran, H., 2001, ApJ, 554, L19
Turner, T.J, George, I.M., Nandra, K., Mushotzky, R.F., 1997, ApJ, 488, 164
Winkler, H., Glass, I.S., van Wyk, F., Marang, F., Jones, J.H.S., Buckley, D.A.H., Sekiguchi, K., 1992, MNRAS, 257, 659
Young, S., Hough, J.H., Efstathiou, A., Wills, B.J., Bailey, J.A., Ward, M.J., Axon, D.J. 1996, MNRAS, 281, 1206



Figure 1: The observed correlations between L$_{25}$ and (a) L$_{\rm HX}$ and (b) L$_{5007}$. The Seyfert 1s are shown as ∘, the Compton thin Seyfert 2s as • and the Compton thick Seyfert 2s as +. 25μm upper limits are shown by the arrows. The solid straight line in each plot is the best fit to the data excluding those galaxies which are Compton thick, and those galaxies in (a) and (b) L$_{\rm HX} \lesssim 10^{7.2}$ L$_\odot$ and L$_{5007} \lesssim 10^{7.3}$ L$_\odot$. The vertical dashed lines show the boundaries between those galaxies which we have classed as AGN dominated, and those in which the host galaxy dominates. Excess 25μm emission can clearly be seen for the galaxies to the left of these lines.



Figure 2: (a) IRAS colours of the sample as a function of L$_{5007}$. The Seyfert 1s and 2s have been merged in this sample. Those with L$_{5007} < 10^{7.3}$ L$_\odot$ are shown as +. These are the galaxies which we class as host galaxy dominated. Those with $10^{7.3} <$L$_{5007} < 10^{8.5}$ L$_\odot$ are shown as △. Those with L$_{5007} > 10^{8.5}$ L$_\odot$ are shown as ⋆. The solid line plotted is the track followed by an increasingly reddened Seyfert 1. The dashed line shows the locus of the starburst/LINER population. The dot-dashed line is the boundary within which galaxies of mixed AGN and starburst excitation should lie. See Dopita et al. (1998) for more details. (b) The compactness of the 10μm emission as a function of the L$_{5007}$. Seyfert 1s are shown as ∘, and Seyfert 2s as •, clearly indicating the lower luminosity objects are more likely to be less compact. Only galaxies with $cz < 5000$kms^{-1} are shown.


(a)

(b)

(c)

Figure 3: Luminosity distribution of the galaxies in which HBLRs are detected (solid line) and not detected (dashed line) as a function of the L_{25} for the (a) Moran et al. (2000) sample, (b) Tran (2001) sample and (c) the L01 sample.